

# Two Steps for Controlling SCN

## 1. Rotate, Rotate and Rotate Crop and Variety Rotations

- Rotate with non-host crops such as corn, wheat or alfalfa.
- If SCN numbers are high, rotate with soybean varieties having different sources of resistance.
- Four genetic sources of resistance currently are available.
- The use of one source of resistance will eventually lead to SCN populations that overcome that resistance.
- Soybean varieties with similar sources of resistance can differ in their ability to suppress reproduction, so it is important to switch varieties.
- Rotate with tolerant or susceptible varieties when SCN numbers are low.

## 2. Relieve Stress

Manage these to avoid compounding damage due to SCN:

- **weeds**
- **water**
- **fertility**
- **herbicide**



**SCN** is the **most destructive pest** of soybeans in the world. **Understanding nematode biology** is key to recognizing **early infestation** and **managing** its **negative economic impact**.

### Example rotation schedule:

<b>Year 0</b>	Identification of SCN
<b>Year 1</b>	Non-host crop (corn, alfalfa, wheat)
<b>Year 2</b>	Adapted resistant variety
<b>Year 3</b>	Non-host crop
<b>Year 4</b>	Susceptible soybean variety if soil analysis shows that SCN levels are below threshold, resistant variety if levels are still high.
<b>Year 5</b>	Repeat rotation sequence

# SCN Key Points

1. Find out if you have SCN, you may have it and not know it.
2. SCN may not cause obvious symptoms.
3. SCN can cause substantial yield loss without causing symptoms.
4. If you have SCN, follow local recommendations.

### For more information, contact:

UW Agronomy Soybean Research and Extension  
1575 Linden Drive  
Madison, WI 53706  
608.262.7975  
[www.coolbean.info](http://www.coolbean.info)

Field Crops Pathology  
[fyi.uwex.edu/fieldcroppathology/soybean](http://fyi.uwex.edu/fieldcroppathology/soybean)

The Soy Report  
[thesoyreport.blogspot.com](http://thesoyreport.blogspot.com)

North Central Soybean Research Program  
[www.ncsrp.com](http://www.ncsrp.com)

### Free SCN Testing Available to Wisconsin Growers

Program funded by WSMB.  
Request up to 4 free kits for your farm:  
[freescntest@mailplus.wisc.edu](mailto:freescntest@mailplus.wisc.edu)


Funded by the soybean checkoff through the Wisconsin Soybean Marketing Board (WSMB)  
[www.wisoybean.org](http://www.wisoybean.org)

# Wisconsin Soybean Cyst Nematode Management

## The Billion Dollar Pest

What you Can't See, Will Hurt You.


## Guide For Wisconsin



## Recognizing a SCN Problem

- Symptoms of SCN infections can range from no visible evidence of plant injury to plant death in certain areas of the field.
- Chlorosis (yellowing) is caused by nitrogen deficiency because of the ability of SCN to inhibit Rhizobium nodule formation.
- Areas are patchy, circular or oval in shape.
- Symptoms associated with SCN damage are similar to other crop production problems such as potassium and nitrogen deficiencies, iron chlorosis, herbicide injury, soil compaction, drought stress and other soybean diseases.
- White or yellow female is the only visible sign of SCN infection, but they may not be present at the time of sampling. Cysts (dead brown females) are not visible in soil.
- White female SCN are seen in the field when the plant begins to flower. While observation of white females will confirm an SCN infestation, it tells you little about the level of infestation. If you dig up roots and don't find a white female, that doesn't mean SCN is absent. The only way to get a reliable diagnosis is through a professional diagnostic laboratory.

White females of SCN on roots are visible to the naked eye, but are still very small (here magnified 32X). Nitrogen-fixing nodules are much larger.



## Soil Sampling for SCN

Soil can be sampled any time for the presence of SCN. A good time to sample soil is in the fall, before the soybean harvest during routine sampling to determine soil fertility.

### Procedure:

1. Use a soil probe or narrow-bladed trowel or shovel. Take cores close to plants at a depth of 6 to 8 inches. Be sure to include plant roots.
2. Submit one sample for each 10-acre field, or for a suspected area within the field. Sample from plants in the margins of suspected areas, and not from their centers. Plants in the center of the affected area have severely stunted root systems that cannot support SCN. Or collect in a zig-zag pattern across the field. Collect from areas of similar soil texture and cropping history. If different crops were grown, or there is markedly different soils within a field, sample separately.
3. Take soil and roots from 12 to 20 cores/10 acres and mix into one sample. Place 1 pt. (2 cups) of mixed soil into a plastic soil sample bag, fasten the open end securely, and write your name on the bag label. Keep the samples out of the sun and don't let them dry out.
4. Mail as soon as possible and early in the week to avoid delays in transit.




## Understanding SCN Soil Test Results

There is a relationship between the number of SCN eggs present at the time of planting and soybean yield. To test a sample for SCN, the soil is added to water and then passed over a sieve to remove the cyst (female) stage of the nematode. The cysts are separated from debris by a centrifugation technique and then crushed to release the eggs. Each cyst may contain 10-250 eggs. Egg counts are used to assess the risk of yield loss due to SCN. SCN results are usually expressed as the number of eggs per volume of soil. Some labs report number of cysts per volume of soil. Egg population density is a better predictor of risk than cyst number. This table will help you understand your SCN lab report.

Risk	Identification of SCN	Potential Yield loss for SCN susceptible variety
None	0 eggs	None
Low	1-500 eggs	0-10% silt or clay soils
Moderate	500-2000 eggs 1-500 eggs	10-20% silt or clay soils 5-20% sandy soils
High	500-5,000 eggs	10-50% all soils
Very High	>5,000 eggs	Very high-expect yield loss for resistant variety

### Fields Testing Negative for SCN

May not remain free since this pest can be introduced in soil, moved by animals, machinery or wind.



This SCN cyst has been broken open to reveal the eggs inside. Eggs can remain viable for years even in the absence of a suitable host.

# The “yield robber” Soybean Cyst Nematode

A disease that can live for many years in the soil, and is relatively **invisible** when looking at an **infested field**.

## Overview

- **Soybean Cyst Nematode (SCN)** was first found in the U.S. in North Carolina in 1954.
- SCN is the most **serious** soybean pest in the U.S.
- SCN causes more than **\$1 billion in soybean yield losses** each year.
- A microscopic roundworm, the nematode occurs in all **major soybean production areas**.
- SCN causes **no specific symptoms** and its effects are often not dramatic.
- Many growers do not know they have a problem until a **severe infestation develops**.
- **Yield losses** due to SCN can be over **50%**.
- Understanding the life cycle of SCN, **routine soil testing** and **proper crop management** can **reduce** the incidence of this **pest**.
- SCN **resistant soybean varieties** will **increase yield** potential of infested fields.